

Landau level bosonization of a 2D electron gas

H. Westfahl Jr.¹, A.H. Castro Neto² and A.O. Caldeira¹

¹*Instituto de Física Gleb Wataghin
Universidade Estadual de Campinas
13083-970, Campinas, SP, Brazil*

²*Department of Physics
University of California
Riverside, CA, 92521, USA*

(February 1, 2008)

In this work we introduce a bosonization scheme for the low energy excitations of a 2D interacting electron gas in the presence of an uniform magnetic field under conditions where a large integral number of Landau levels are filled. We give an explicit construction for the electron operator in terms of the bosons. We show that the elementary neutral excitations, known as the *magnetic excitons* or *magnetoplasma modes*, can be described within a bosonic language and that it provides a quadratic bosonic Hamiltonian for the interacting electron system which can be easily diagonalized.

PACS: 73.40.-c, 73.40.Hm, 73.20.Mf, 71.27.+a

Bosonization of Fermi fields has been an important non-perturbative tool for finding the solutions to problems involving many interacting fermions. The pioneering attempts by Bloch¹ on sound waves in dense 1D systems, which were later extended and given an intuitive foundation by Tomonaga², called attention to the conceptual and operational simplicity that results if all excitations of a fermionic system can be described within a bosonic language³⁻⁵. Nowadays the 1D bosonization is a mature field⁶. Bosonization in higher dimensions was pioneered by Luther⁷ and extended by Haldane a few years ago⁸. More recently this approach has been applied in the context of strongly correlated systems where the possibility of non-conventional ground states seeks for a non-perturbative method of study⁹⁻¹¹.

In the context of correlated systems, one of the most interesting phenomenon in condensed matter physics is the quantum Hall effect¹²⁻¹⁴ either in its integer (IQHE) or fractional (FQHE) regimes. Despite the fact that there is no thermodynamic distinction between this two phenomena¹³, they have quite different microscopic origins. The IQHE can be understood from the properties of non-interacting electrons whereas the understanding of the FQHE necessarily requires a treatment of the electron-electron interaction. Consequently, almost all the efforts to study correlations in a 2D electron gas in the presence of a magnetic field, B , were directed toward the study of the strong magnetic field case where the FQHE takes place. In this case the Landau level spacing, the cyclotron energy $\hbar\omega_c = \hbar eB/(mc)$, can be much greater than the typical Coulomb energy, which is of the order of e^2/ℓ (where $\ell = \sqrt{\hbar/(eB)}$ is the magnetic length), and therefore the system can be described only in terms of the first Landau level and the mixing with the other Landau levels can be neglected or taken into account perturbatively. In a weak magnetic field¹⁵ this is not the case and the typical Coulomb energy can exceed the cyclotron energy resulting in a mixing between the Landau levels. Some attempts were made to develop a

method for studying the mixing caused by these correlation effects starting from a Fermi Liquid theory^{16,17} or perturbative Green's function method^{15,18,19}.

In this paper we start from a Landau level description of the system and introduce a non-perturbative bosonization scheme for the 2D interacting fermion gas subject to a perpendicular uniform magnetic field under the condition that there is a large integer number of filled Landau levels. In this case, the excitations are electron-hole pairs often called magnetic excitons and as we will show can be described within a bosonic language.

Recent experiments in tunneling between parallel 2D electron gases in applied magnetic fields²⁰ have revealed the existence of a gap in the tunneling density of states. In this case it is established that correlations have major hole in the tunneling conductance and we expect that the method presented here can be applied to such problems where the correlations between the electrons in a magnetic field are so important.

We start with the Hamiltonian of a spinless fermion gas in a plane perpendicular to an uniform magnetic field

$$H_o = \frac{1}{2m} \int d^2\mathbf{r} \Psi^\dagger(\mathbf{r}) \left(\mathbf{p} - \frac{e}{c} \mathbf{A}(\mathbf{r}) \right)^2 \Psi(\mathbf{r}),$$

where $\Psi(\mathbf{r})$ is the fermion field operator (for the time being we do not discuss the spin of the electron).

Rewriting $\Psi(\mathbf{r})$ in a Landau level basis (symmetric gauge) the Hamiltonian H_o becomes diagonal

$$H_o = \hbar\omega_c \sum_{n,m=0}^{\infty} (n+1/2) c_{n,m}^+ c_{n,m},$$

where $c_{n,m}^+$ is the fermion operator that creates a particle in a Landau level n with the guiding center at a distance $\sqrt{2m+1}\ell$ from the origin of the coordinate system. Therefore the m degeneracy is related to the location of the position of the guiding center of the cyclotron orbit.

The non-interacting ground state is generated by uniformly filling ν Landau levels (ν integer) of each guiding center labeled by the quantum number m

$$|G\rangle = \prod_{m=0}^{N_\phi} \prod_{n=0}^{\nu-1} c_{n,m}^+ |0\rangle,$$

where $N_\phi = \frac{S}{2\pi\ell^2}$ is the number of flux quanta. The neutral excitations in the non-interacting system can be generated by creating a hole in a Landau level $p \leq \nu - 1$ at the guiding center m and an electron in a level $n + p \geq \nu - 1$ at the guiding center m' .

We note, however, that it is not possible to define eigenstates of the guiding center $\vec{R}_{0i}$ for the cyclotronic motion of a single charged particle because of the non-commutativity of the components of the position operators. Nevertheless we can define a vector $\vec{R}_0 = \vec{R}_{0e} + \vec{R}_{0h}$, related to the center of mass of the guiding centers of the electron and the hole excited in the system, whose components x_0 and y_0 commutes with each other and therefore have simultaneous eigenstates. It is also possible to define a momentum operator $\vec{P}_0$, canonically conjugated to $\vec{R}_0$, which is related to the momentum of the center of mass of the guiding centers. The eigenstates of this operator can be generated by a superposition of the above described neutral excitations in different guiding centers and they form a complete basis in the space of the neutral excitations known as *magnetic excitons*¹⁸. In operator form they can be described as,

$$e_{n,p}^\dagger(\mathbf{k}) |G\rangle = e^{-\frac{(\kappa\ell)^2}{4}} \sum_{m',m=0}^{\infty} G_{m',m}(\ell k) c_{n+p,m'}^+ c_{p,m} |G\rangle,$$

where $k = k_x + ik_y$, $\kappa = |k|$ and $G_{m',m}$ is given by¹⁴

$$G_{n+p,p}(k) = \sqrt{\frac{p!}{(n+p)!}} \left(\frac{-ik}{\sqrt{2}}\right)^n L_p^n\left(\frac{\kappa^2}{2}\right)$$

with L_p^n being generalized Laguerre polynomials.

The density fluctuations of the system can be expressed in terms of superpositions $a_n^\dagger(\mathbf{k})$ of these *magnetic excitons*

$$\rho(\mathbf{k}) - \rho_0 = \sum_{n=1}^{\infty} \{a_n^\dagger(\mathbf{k}) + a_n(-\mathbf{k})\},$$

where

$$a_n^\dagger(\mathbf{k}) = e^{-\frac{(\kappa\ell)^2}{4}} \sum_{p=0}^{\infty} G_{n+p,p}(\ell k^*) e_{n,p}^\dagger(\mathbf{k}) \quad (1)$$

can be interpreted as *magnetic plasmons* and $\rho_0 = \frac{\nu}{2\pi\ell^2}$ is the electronic average density in the system.

In order to bosonize this problem we proceed exactly like in the Luttinger model and enlarge the Hilbert space of the fermionic system by including filled energy eigenstates of negative values of the quantum number n . This construction is the analogue of the sea of infinite negative

energy of the Luttinger model (the Tomonaga construction can also be obtained in an analogous way by making the Hilbert space finite²¹). This vacuum does not concern us because we are studying the *low energy excitation* spectrum of the system. It is clear from this construction that the bosons only make a faithful representation of the electronic system if $\nu \gg 1$. In the same way the bosonization of the electron gas is only possible when the system is very dense (large Fermi momentum). By doing that one can show that the bosonic operators defined in (1) obey the *exact* commutation relations,

$$\begin{aligned} [a_n(\mathbf{k}), a_{n'}^\dagger(\mathbf{k}')] &= \delta_{n,n'} \delta_{\mathbf{k},\mathbf{k}'} N_\phi \mathcal{G}_n(k\ell) \\ [a_n(\mathbf{k}), a_{n'}(\mathbf{k}')] &= 0, \end{aligned}$$

where

$$\mathcal{G}_n(x) = n J_n^2(k R_c),$$

J_n is a Bessel function of the first kind and $R_c = \sqrt{2\nu}\ell$ is the size radius of the last occupied “orbit”. We can proceed further and define the canonical bosonic operators

$$b_n(\mathbf{k}) = \frac{1}{\sqrt{N_\phi \mathcal{G}_n(k\ell)}} a_n(\mathbf{k})$$

and recover the standard bosonic commutation relations.

It is easy to show that in terms of these bosonic degrees of freedom the free Hamiltonian can be rewritten, apart from a background energy, as

$$H_o = \hbar\omega_c \sum_{\mathbf{k}} \sum_{n=1}^{\infty} n b_n^\dagger(\mathbf{k}) b_n(\mathbf{k}) \quad (2)$$

which describes the energy of an assembly of non-interacting bosons.

So far we have succeeded in expressing the neutral excitations of the system of electrons in terms of bosons. In order to complete our scheme one needs to express the electron operator in terms of bosons. This is done by transforming from the momentum representation which is the natural representation for bosons to the $|nm\rangle$ representation which is the appropriate one for fermions in a magnetic field. This should be contrasted with the case of the electron gas where the momentum is a good quantum number for the electron and the particle-hole excitation. It means that we must construct an operator that creates an electron in a Landau level n and guiding center m as a function of the density fluctuations within this guiding center. This transformation involves a linear superposition of the magnetic excitons at different momentum states. This is done by considering the operator

$$\begin{aligned} a_{n,m}^\dagger &= \frac{1}{\sqrt{n}} \sum_{p=-\infty}^{\infty} c_{n+p,m}^\dagger c_{p,m} \\ &= \int \frac{d^2\mathbf{k}}{2\pi} e^{-\frac{\kappa^2}{4}} G_{m,m}(k) b_n^\dagger(\mathbf{k}). \end{aligned}$$

Differently from the operator $b_n^\dagger(\mathbf{k})$ that creates density fluctuations with a definite momentum $\mathbf{k}$, the operator $a_{n,m}^\dagger$ generates density fluctuations on a definite

guiding center. This operator also obeys exact bosonic commutation relations. This is completely analogous to Luther's^{7,8} construction of the Fermi liquid in 2D and 3D as many 1D fermi liquids, one for each direction normal to the Fermi surface. But contrary to that construction here we do not need to “smear” the operators close to the Fermi surface. The discrete nature of the Landau states does this smearing naturally. Here we have also reduced the 2D problem of electrons in a magnetic field to N_ϕ 1D *chiral* problems, one for each guiding center labeled by m .

In order to complete the bosonic description of the fermion operator it will be necessary to define an operator that creates a fermion in a “phase” state associated with this circular motion. This is done by rewriting the original fermion operator as,

$$c_m(\theta) = \frac{1}{\sqrt{2\pi}} \sum_{n=-\infty}^{\infty} e^{-in\theta} c_{n,m},$$

where we have used the enlarged Hilbert space and it is easy to show that

$$\{c_m^\dagger(\theta'), c_m(\theta)\} = \delta_{m',m} \delta(\theta' - \theta) \quad (3)$$

$$\{c_m(\theta'), c_m(\theta)\} = 0. \quad (4)$$

The operator $c_m^\dagger(\theta)$ creates an electron about the guiding center given by the quantum number m and at the phase θ of the cyclotronic motion. This is the operator that can be properly bosonized. Indeed the Mandelstam form^{4,5} of the fermion operator is given by

$$c_m^\dagger(\theta) = \frac{1}{\sqrt{2\pi}} e^{-i\nu\theta} e^{i\phi_m^\dagger(\theta)} U_m e^{i\phi_m(\theta)},$$

with

$$\phi_m(\theta) = \frac{N_m \theta}{2} - i \lim_{\alpha \rightarrow 0} \sum_{n=1}^{\infty} e^{-\frac{n\alpha}{2}} \frac{e^{-in\theta}}{\sqrt{n}} a_{n,m}$$

and

$$U_m = e^{i\pi \sum_{s=0}^{m-1} N_s} e^{i\Theta_m}$$

where the Hermitian operators

$$N_m = \sum_{p=-\infty}^{\infty} (c_{p,m}^\dagger c_{p,m} - c_{p,m}^\dagger c_{p,m} >)$$

and Θ_m are canonically conjugated ($[N_m, \Theta_{m'}] = i\delta_{m'm}$) such that the unitary operators U_m anticommute.

In this representation, it can be shown that the fermionic operators obey the desired anticommutation relations (3) and (4) and it recovers the long time asymptotic behavior of the non-interacting propagator between two different phase states which is given by

$$\begin{aligned} K(\theta, t) &= \langle c_{m'}^\dagger(\theta, t) c_m(0, 0) \rangle \\ &= \delta_{m',m} \frac{e^{-i\nu\theta}}{1 - e^{-i(\theta - \omega_c t)}} \end{aligned}$$

and can be related to the space-time propagator by writing the operator $\Psi(\mathbf{r})$ in terms of the $c_m(\theta)$ ²¹. That completes the construction of the model. Now we have an *operator* relation between electrons in a magnetic field and bosons.

Our next step is to include the interaction between the electrons within this framework. In an uniform neutralizing background the total Hamiltonian can be written as

$$H = H_o + H_I$$

where H_o is given by (2) and

$$\begin{aligned} H_I &= \frac{1}{\ell^2} \sum_{\mathbf{k}} V(k) \sum_{n,n'=1}^{\infty} \sqrt{\mathcal{G}_n(k\ell) \mathcal{G}_{n'}(k\ell)} \\ &\quad \{b_n^\dagger(\mathbf{k}) + b_n(-\mathbf{k})\} \{b_{n'}^\dagger(-\mathbf{k}) + b_{n'}(\mathbf{k})\} \end{aligned}$$

is a density-density interaction between the fermions which is written in bosonic language. $V(k)$ is the Fourier transform of the interacting potential.

In order to diagonalize H we will adopt a generalized Bogoliubov transformation¹⁰ with real coefficients $\mu_{nl}(\mathbf{k})$ and $\vartheta_{nl}(\mathbf{k})$ which is given by

$$b_n(\mathbf{k}) = \sum_{l=1}^{\infty} \left(\mu_{nl}(\mathbf{k}) \beta_l(\mathbf{k}) + \vartheta_{nl}(\mathbf{k}) \beta_l^\dagger(-\mathbf{k}) \right),$$

where $\beta_l(\mathbf{k})$ are the new bosonic operators. This canonical transformation yields

$$H = \hbar\omega_c \sum_{\mathbf{k}} \sum_{l=1}^{\infty} \Omega_l(\mathbf{k}) \beta_l^\dagger(\mathbf{k}) \beta_l(\mathbf{k})$$

where the new eigenfrequencies $\Omega_l(\mathbf{k})$ are determined by the equation

$$1 = \left(\frac{V(k)}{\hbar\omega_c \ell^2} \right) \sum_{n=1}^{\infty} \frac{\mathcal{G}_n(k\ell)n}{(\Omega_l(\mathbf{k})^2 - n^2)}.$$

The above equation has the same form as the one obtained for the normal modes in the integer quantum Hall effect in the RPA approximation for a Coulomb potential, $V(\kappa) = 2\pi e^2/\kappa$ ¹⁸. Moreover, the same equation is obtained in the fractional quantum Hall effect for composite fermions¹⁹. Now we have shown that as far as the bosonization construction is valid, the equation which defines the eigenvalues holds for any value of the magnetic field. This is a consequence of the fact that bosonization fulfills the Ward identities. Thus, exactly like in the Fermi gas, the limit of validity of bosonization goes beyond what is expected.

As a last comment we would like to mention that the method developed in this letter can be applied to the fractional Hall effect if we use the Chern-Simons theory to map the original electrons into the composite fermions. In this case, as it is well established, the fractional quantum Hall effect is mapped into the integer quantum Hall effect of composite fermions. The condition that $\nu \gg 1$

becomes equivalent to say that we are looking at fillings close to $1/2$ ²². The nature of the ground state of this system has recently been the source of much controversy and we believe that our methods can provide non-perturbative answers for the structure of the ground state of this problem²¹.

In conclusion, we have proposed a new method of bosonization for 2D fermions subject to a uniform magnetic field by directly working in a Landau basis. We propose an analogue of the Luttinger model which can be bosonized and provides an operator relation between electrons and bosons in the limit of $\nu \gg 1$. This construction is essentially non-perturbative and provides new insights into the problem of electrons interacting in the presence of magnetic fields. We show that bosons are collective excitations of the system which are related with a collection of N_ϕ 1D Luttinger liquids associated with each guiding center. We also obtain the dispersion of the collective modes of the system and discuss the applicability of the method to the integer and fractional quantum Hall effects. We expect our method will enlarge the understanding of the physics of strongly correlated systems in the presence of magnetic fields.

H.W. and A.O.C. kindly acknowledge respectively the full and partial supports from the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). We are also grateful to E.H. Fradkin, F.D.M. Haldane and D. V. Khvashchenko for very useful discussions.

ences therein.

- ¹⁴ A.H. MacDonald, *The Quantum Hall Effect: A Perspective*, Kluwer Academic Dordrecht, 1989.
- ¹⁵ A.H. MacDonald, Phys. Rev. B **30**, 4392 (1984).
- ¹⁶ I.L. Aleiner and L.I. Glazman, Phys.Rev.B, **52**, 11296, (1995).
- ¹⁷ L.S. Levitov and A.V. Shytov, cond. mat. /9507058.
- ¹⁸ C. Kallin and B.I. Halperin, Phys. Rev.B, **30**, 5655, (1984).
- ¹⁹ A.Lopez and E.Fradkin, Phys. Rev. B **44**, 5246 (1991); Phys. Rev. Lett. **69**, 2126 (1992); Nucl. Phys. B **33 C**, 67 (1993).
- ²⁰ J.P.Eisenstein, L.N. Pfeiffer and K.W. West, Phys. Rev. Lett. **74**, 1419 (1995).
- ²¹ H.Westfahl Jr., A.H. Castro Neto and A.O. Caldeira, in preparation.
- ²² B. I. Halperin, P. A. Lee and N. Read, Phys. Rev.B **47**, 7312 (1993).

-
- ¹ F. Bloch, Z. Phys. **81**, 363 (1933).
 - ² S. Tomonaga, Prog. Theor. Phys. (Kyoto), **5**, 544, (1950).
 - ³ W. Thirring, Ann. Phys. (N. Y.) **3**, 91 (1958); J.M.Luttinger, J.Math.Phys. **4**, 1154, (1963); D. Mattis and E. Lieb, J. Math. Phys. **6**, 304 (1965); A. Luther, Phys. Rev. B **14**, 2153 (1975); S. Coleman, Phys. Rev. D. **11**, 2088 (1975).
 - ⁴ S. Mandelstam, Phys. Rev. D **11**, 3026 (1975).
 - ⁵ F.D.M. Haldane, J. Phys. C **14**, 2585 (1981).
 - ⁶ For a recent review on this subject see, J. Voit, Rep.Prog.Phys. **58**, 977 (1995) and references therein.
 - ⁷ A.Luther, Phys.Rev.B, **19**, 320 (1979).
 - ⁸ F.D.M. Haldane, Varenna Lectures (1992) and Helv. Phys. Acta. **65**, 152 (1992).
 - ⁹ A. Houghton and B. Marston, Phys. Rev. B **48**, 7790 (1993); H.J. Kwon, A. Houghton and B. Marston, Phys. Rev. B **52**, 8002 (1995).
 - ¹⁰ A.H. Castro Neto and Eduardo Fradkin, Phys. Rev. Lett. **72**, 1393 (1993); Phys. Rev. B, **51**, 4048 (1995).
 - ¹¹ P. Kopietz and G.E. Castilla, Phys.Rev.Lett. **76**, 4777 (1996); P.Kopietz, J. Hermisson and K. Schonhammer, Phys.Rev.B **52**, 10877 (1995).
 - ¹² R.E. Prange and S.M. Girvin eds., *The Quantum Hall Effect*, 2nd ed. (Springer-Verlag, New York, 1990).
 - ¹³ A.Karlhede, S.A. Kivelson, and S.L. Sondhi, *The Quantum Hall Effect : The article*, Lectures pres. at the 9th Jerusalem Winter School on Theor. Phys. (1992) and ref-